



# International Journal of Mechanical and Thermal Engineering

E-ISSN: 2707-8051

P-ISSN: 2707-8043

IJMTE 2023; 4(2): 60-68

Received: 24-06-2023

Accepted: 26-07-2023

**Stanislav Ordin**

The Russian Academy of  
Sciences, Moscow, Russia

## Non-Schroedinger Orbitals

**Stanislav Ordin**

### Abstract

Schrödinger's "Searchlight" allowed the atom to be illuminated and cast its shadow on the "Observer's Screen". But the main problem of the modern "Quantum Theory" is precisely that it mistook the shadows from the "Elephant" for the "Elephant" itself, for Reality. This led to the stagnation of the "Theory" and, with an increase in the mass of atoms, to a catastrophic discrepancy between the allowed calculated electron energy levels and the experimental Ionization Potentials. But the inclusion of "normal illumination" - Understanding gives a "holographic image" - a real Quantum Theory, which can be correctly built on the path laid by Planck-de Broglie-Einstein-Heisenberg-Bohm. Quantization supplements with a new Invariant and, strictly mathematically according to Planck, expands and complements the Classics, and does not deny it. In this case, the resonant orbits of electrons in atoms - Electron Orbitals correspond to the resonant de Broglie waves and Pontryagin's Dualism of Functional Sets translates Schrödinger's Uncertainty Principle into the Principle of Certainty = Observability.

**Keywords:** Classical orbit, Ionization Potential, Planck resonance, de Broglie waves

### Introduction

"Quantum theory explains a lot, but in fact it does not bring us one step closer to the secrets of the Old Man, in any case, I am convinced that He does not play dice" Albert Einstein

Introduced by Bohr in the planetary model of the Rutherford atom, Stationary Orbits of electrons <sup>[1, 2]</sup>, he himself, using quantum, without realizing it, pushed his own Orbits, moving his founders - Planck and Einstein - away from the Fundamentals of Quantization. Fascinated by the mysticism of mathematical solutions to the Schrödinger equation, Bohr replaced his three-dimensional orbits, due to his poor understanding of mathematics <sup>[3]</sup>, with interpretations of the imaginary Schrödinger "wave function". So, this mysticism was included in their canonized definition itself: *Atomic orbital (electron orbital) is a one-electron wave function,  $\psi$ , obtained by solving the Schrödinger equation for a given atom; is given by the principal  $n$ , orbital  $l$  and magnetic  $m$  - quantum numbers <sup>[4]</sup>.*

But this definition of Orbital included in the primers, strictly speaking, is not its definition, but is its redefinition, moreover, looped, like a snake devouring its own tail. So the entire explanation of Chemistry by "Quantum Mechanics" is based on an initially false definition <sup>[5]</sup>. Indeed, this "definition" does not appeal to known measurable waves - to the Heaviside-Maxwell electromagnetic wave quantized by Planck or to the acoustic wave quantized by Einstein, and not to the de Broglie matter wave, albeit somewhat mystical (for now), but experimentally observable, quantization which, as will be shown, gives stationary - resonant orbits of electrons. No, this "definition" appeals to a fundamentally unobservable "wave function". Those. Without losing the underlying "meaning," this "definition" (with the same success and with the same errors) could be rewritten the other way around: "The wave function is a one-electron atomic orbital (electron orbital)." But the physicists who gathered in Copenhagen over a bottle of cheap port came up with "their own" special name for the "wave function", in order to somehow tie it to reality: probability density. But they didn't tie it tightly - Schrödinger worked out that it can be not only non-zero, but maximum inside the nucleus! There are other reasons indicating that the "wave function" has nothing to do with the probability density <sup>[6]</sup> and in general, it turned out the same way as when referring to the Principle of Causality simply disguises the fact that the reason for something is not understood.

**Corresponding Author:**

**Stanislav Ordin**


The Russian Academy of  
Sciences, Moscow, Russia

# Quantities, Units and Symbols in Physical Chemistry

Prepared for publication by

IAN MILLS    TOMISLAV CVITAŠ  
KLAUS HOMANN    NIKOLA KALLAY  
KOZO KUCHITSU

SECOND EDITION



BLACKWELL SCIENCE

http://goldbook.iupac.org/A00500.html

© 1993 International Union of Pure and Applied Chemistry and published for them by Blackwell Science Ltd

Editorial Offices:  
Osney Mead, Oxford OX2 0EL  
25 John Street, London WC1N 2BL  
23 Ainslie Place, Edinburgh EH3 6AJ  
359 Main Street, Malden  
MA 02148 5018, USA  
54 University Street, Carlton  
Victoria 3053, Australia  
10, rue Casimir Delavigne  
75006 Paris, France

Other Editorial Office:  
Blackwell Wissenschafts-Verlag GmbH  
Kurfürstendamm 57  
10707 Berlin, Germany

Blackwell Science KK  
MG Kodomacho Building  
7-10 Kodomacho Nihombashi  
Chuo-ku, Tokyo 104, Japan

The right of the Author to be identified as the Author of this Work has been asserted in accordance with the Copyright, Designs and Patents Act 1988.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, except as permitted by the UK Copyright, Designs and Patents Act 1988, without the prior permission of the copyright owner.

First published 1988  
Reprinted 1988  
Reprinted as paperback 1989  
Russian translation 1988  
Hungarian translation 1990  
Indian reprint edition 1990  
Japanese translation 1991  
Second edition 1993  
Reprinted 1995, 1996, 1998

Set by Macmillan India Ltd  
Printed and bound in Great Britain at the University Press, Cambridge

The Blackwell Science logo is a trade mark of Blackwell Science Ltd, registered at the United Kingdom Trade Marks Registry

**DISTRIBUTORS**

Marston Book Services Ltd  
PO Box 269  
Abingdon  
Oxon OX14 4YN  
(Orders: Tel: 01235 465500  
Fax: 01235 465555)

USA and Canada  
CRC Press, Inc.  
2000 Corporate Blvd, NW  
Boca Raton  
Florida 33431

Australia  
Blackwell Science Pty Ltd  
54 University Street  
Carlton, Victoria 3053  
(Orders: Tel: 3 9347 0300  
Fax: 3 9347 5001)

A catalogue record for this title is available from the British Library

ISBN 0-632-03583-8

Library of Congress  
Cataloging in Publication Data

Quantities, units and symbols in physical chemistry/  
prepared for publication by  
Ian Mills... [et al.]—2nd ed.  
p. cm.  
At head of title: International Union of Pure and Applied Chemistry  
International Union of Pure and Applied Chemistry, Physical Chemistry Division,  
Commission on Physicochemical Symbols, Terminology, and Units—P. facing t.p.  
Includes bibliographical references and index.  
ISBN 0-632-03583-8  
I. Chemistry, Physical and theoretical—Notation.  
II. Mills, Ian (Ian M.)  
III. International Union of Pure and Applied Chemistry.  
Commission on Physicochemical Symbols, Terminology, and Units.  
QD451.5.Q36 1993  
541.3'014—dc20

If we talk about probability density, then we need to add - What! Thus, the probability density of the kinetic energy of molecules in a gas can also be used to describe a sound wave. Probability densities can be compared not only with electromagnetic waves, but also with the static Coulomb-Newton Laws [7], if, in accordance with the principle of Logarithmic Relativity, we associate with it the sub particles that form these fields and the substructure of the corresponding fields, even if not yet measured by us. And for electron orbitals, if we talk about the probability density of the distribution of an electron in an orbit, then we must honestly say - What! What is the parameter of electrons? And don't hide behind the words Unknown what, when calculations give the maximum "wave" function of the electron inside the nucleus.

So, at the dawn of the last century, the times of Newton, when the Physicist and the Mathematician were in the same "bottle" of Reality, have passed. And the developers of "Quantum Theory" hid their misunderstanding of some aspects of Classical Physics [8], multiplied by their misunderstanding of Planck-Einstein Quantum [9], behind an imaginary unit from mathematics. Heavieside physics, which was also not fully understood, adding "unobservable zero-point oscillations" to the picture for "completeness". And, thus, they turned the entire "Quantum Mechanics" into a schizophrenic Mind Game divorced from Reality. And so it took a hundred years for the intoxicating "charm" of mysticism in the Elementary Harmonic Oscillator, which, in fact, was not even strictly analyzed [10], to pass and a severe hangover set in with "Black Holes" and "Particles of God". If you ate to drink, it would still be better to have good cognac. And with a "sober head" it becomes obvious that the primary quantization of spatial waves in extended media is done correctly - according to the Principles laid down by

Planck and Einstein. This determined the progress of the Quantum Theory of Solid State. But the "quantum" transition to local oscillations has not been thoroughly worked out in its fundamentals and it, even in the simplest case of the Ideal Harmonic Oscillator, gives mystical ("immeasurable") Zero oscillations. So "Secondary Quantization" and Atomic Physics and, after them, Quantum Field Theory for a hundred years were not engaged in correcting Basic Models [11-13], but, in fact, sucking out corrections that exceeded what the corrections correct.

That is why the actually observed dualism [14], even Planck's light quanta, and even more so de Broglie's particles, were actually attributed to mysticism [15, 16], which does not need to be understood, but rather accepted.

Whereas Nature "invented" the first coherent waves arising due to Quantization, into which packets of de Broglie waves filling a particle are transformed, in the form of electron orbitals long before people mastered radio waves and invented a maser-laser, where the property of Bose particles is manifested be in one state without limiting their number. And the manifestation of both the Pauli principle and the same property of Bose particles can be easily demonstrated using the example of electron orbitals, but not Schrödinger's at all, but the original ones - Bohr's. But first we will show that the opposition of classical Bohr orbits to quantum physics is far-fetched, simply dictated by the struggle for the priority of the "invention" of quantum. Bohr or Planck? Half a century later, Pontryagin's Dualism of Functional Sets "reconciled" them. But the canonization of Schrödinger's "shadows", despite the catastrophic discrepancies between calculations based on "shadows" and experiment, prevented us from noticing the difference

between the shadows of the “Elephant” and the Real Elephant for another half a century.

**Classic Orbital**

It follows from Classical Physics that in the hydrogen atom the centrifugal force acting on the electron is equal to the force of the Coulomb attraction of the electron and the nucleus. So the force acting on an electron in orbit is the centrifugal force

$$F_C = m\omega^2 r = m \frac{v^2}{r} \tag{1}$$


And the force of the Coulomb attraction of the electron to the hydrogen nucleus

$$F_{11} = \frac{e^2}{4\pi\epsilon_0 r^2} \tag{2}$$

From the equality of these forces acting on an electron in orbit, we obtain the dependence of the electron velocity in orbit on its radius (Fig. 1)

$$m \frac{v^2}{r} = \frac{e^2}{4\pi\epsilon_0 r^2} \Rightarrow v_1^2 = \frac{e^2}{4\pi\epsilon_0 r_1 m} \Rightarrow v_1 = \pm \frac{e}{2\sqrt{\pi\epsilon_0 m}} \frac{1}{r_1^{1/2}} \tag{3}$$

$$\frac{v_1}{c} = \pm \frac{e}{2c\sqrt{r_1\pi\epsilon_0 m}} \frac{1}{(r/r_1)^{1/2}} = \pm 0.007291686044471947 \frac{1}{(r/r_1)^{1/2}} \tag{4}$$



**Fig 1:** The dependence of the relative (reduced to the speed of light) velocity of the steady motion of an electron in the orbit of a hydrogen atom on the relative radius of the electron orbit, reduced to its experimental radius (blue line)

$e = -1.60 \cdot 10^{-19}$  C,  
 $m = 9.11 \cdot 10^{-31}$  kg,  
 $\epsilon_0 \approx 8.85 \cdot 10^{-12}$  m<sup>-3</sup> kg<sup>-1</sup> s<sup>4</sup> A<sup>2</sup>,  
 $r_1 = 53 \cdot 10^{-12}$  m

The resulting dependence of the electron speed, as well as the speed reduced to the speed of light, are determined not only by world constants, but also by the absolute value of the orbit radius. So, using the tabular value of the size of the hydrogen atom as the diameter of the orbit, we get the red line shown in Fig. 1, a quite reasonable value for the speed of stable rotation of an electron in orbit (the reduced radius is used on the graph). But relatively weak influences can take the electron away from the initially given radius of the classical orbit to neighboring ones.


In this orbit, the potential (Coulomb) energy of an electron, if we set it to zero for an electron at infinity, is equal to

$$U_{11} = \frac{e^2}{4\pi\epsilon_0 r_1} \tag{5}$$

With the speed obtained in formula 3, the kinetic energy of an electron in this orbit is equal to half the potential:

$$E_c = m \frac{v_1^2}{2} = m \frac{\left( \frac{e}{2\sqrt{\pi\epsilon_0 m}} \frac{1}{r_1^{1/2}} \right)^2}{2} = \frac{e^2}{8\pi r_1 \epsilon_0} = \frac{U_{11}}{2} = I_1 \approx 13.55 eV \tag{6}$$

And, of course, is also equal, as shown in Fig. 2, to the energy of the electron's exit into vacuum.



**Fig 2:** Electron energy level diagram on a linear scale (left) and on a logarithmic scale (right) the edge of the potential Coulomb well (red line) with exponentially decreasing equipotentials with the electron energy level superimposed on them (green line)


At the same time, the purely classically obtained numerical value of the electron work function (6), which is in good agreement with the tabular value of the Ionization Potential for the hydrogen atom - 13.595 eV.

The whole question is: Why did the electron "choose" from a continuous series of orbits exactly the radius of rotation that corresponds to the Ionization Potential of hydrogen? Although it is likely that the "choice" of this "quantum" radius was simply hidden behind the classical calculation, based on the most experimentally determined value of the Ionization Potential.

So, in the classical representations for the hydrogen atom, as well as in space calculations, we have a continuous series of orbits, with an easy transition from one to another. And the first cosmic velocity for the Earth's satellites is calculated simply on the basis of their minimum radius, equal to the radius of the earth.

But the "Cosmic" velocities of an electron can be tried to be calculated in the model of particle rotation around an infinitely small point. In the simplest particular case - at a fixed speed of rotation of an electron around the nucleus (which, for clarity, we assume equal to unity), the trajectory of its movement relative to the nucleus is described by a modified Archimedes spiral, the radius of which is determined by the ratio of its "start from the nucleus" speed and the total radial acceleration: the balance of the centrifugal forces and forces of the Coulomb attraction of the electron to the nucleus and something else, which gives the binding of the calculated values to the experimentally observed values of the Ionization Potential in the resonant orbit:

$$r(\theta) = v_0 - \frac{a\theta^2}{2} \tag{7}$$



**Fig 3:** The trajectories of an electron that "started from the nucleus" at different speeds

As can be seen from Figure 3, the electron at low speeds will return to the center of rotation and (if it can) skip it. It enters a stationary circular orbit at an initial speed equal to the speed of movement along the orbit, which is its "first cosmic speed". At high starting speeds, the electron also goes into a stationary, but elliptical orbit. And when the "second/third" cosmic velocity is reached, the electron becomes free-it goes to infinity and no longer returns to the nucleus.

We will not carry out this model calculation in detail, since the model with an infinitely small rotation point is idealized and, in accordance with the Principle of Relativity, must also take into account the limitation of the electron speed by the speed of light, while the speed obtained above on the basis of classical concepts is far from the speed of light. But the obtained value for the speed of an electron in a stationary circular orbit (Fig 1.) is quite real and is "the first space velocity for a hydrogen electron. So the reason for the "quantization" of orbits proposed by Bohr - the stabilization of only a few "selected" orbits from the entire series of orbits - must be sought in something else, and not in the limitation of speed by the Principle of Relativity.

But the value of the speed in orbit was itself obtained from its radius, which was assumed to be equal to the radius of the hydrogen atom (which, naturally, is a tautology) and, as will be shown below, the calculation path indicated by Planck gives, even without "Schrödinger's shadows," not the abstract, which, as shown in the work, leads to a catastrophic discrepancy between the calculated allowed energy levels and the First Ionization Potentials with increasing atomic mass [17], but the real the value of the "Selected" radius, consistent, as will be shown later, with the experimental values of the Ionization Potentials and, thus, Bohr's orbital model of the atom is actually restored!

### Quantum addition to the classics

From the equality of the forces acting on an electron in orbit, in the classical consideration, similarly to formula 3 (or simply by dividing formula 3 by the radius), we can also obtain the dependence of the frequency of its rotation around the nucleus on the radius

$$m\omega^2 r = \frac{e^2}{4\pi\epsilon_0 r^2} \Rightarrow \omega_1^2 = \frac{e^2}{4\pi\epsilon_0 r_1^3 m} \Rightarrow \omega_1 = \frac{e}{2\sqrt{\pi\epsilon_0 m}} \frac{1}{r_1^{3/2}} \quad (8)$$

And so, the classical consideration gives one functional relationship of either the electron velocity in the orbit, or the rotation frequency with its radius, using the experimental value of which, according to Bohr, we simply postulated their separation from the series as stationary.

Now we use the Planck equation and the Planck constant to relate to the frequency of light equal to the energy of the Ionization Potential, which, according to formula 6, is equal to the kinetic energy of an electron in orbit

$$I_1 = \hbar\omega_{phl} = E_C = m \frac{v_1^2}{2} = m \frac{\omega_1^2 r_1^2}{2} \quad (9)$$

This equation complements the classical ones and, thus, gives an additional connection between the rotation frequency and the radius of the orbit, if the light quantum correlates with the rotation frequency of the electron in orbit and, thus, if we assume this correlation of frequencies, then

it is possible to obtain a single value of both the frequency and the radius of the classical orbit.

In particular, if we assume that these frequencies are equal, then from (9) we obtain, in addition to equation (8), one more equation relating the frequency to the radius.

$$\omega_1 = \frac{2\hbar}{mr_1^2} \quad (10)$$

And equating the frequencies, we get one equation with one unknown radius:

$$\frac{2\hbar}{mr_1^2} = \frac{e}{2\sqrt{\pi\epsilon_0 m}} \frac{1}{r_1^{3/2}} \quad (11)$$

The solution of the resulting equation gives us the only value of the radius of the RESONANT orbit of an electron in the hydrogen atom

$$r_1 = \frac{16\pi\hbar^2\epsilon_0}{e^2 m} = 212.14\text{pm} \quad (12)$$

So, the World Constants, supplemented by Planck's constant, uniquely DETERMINE the radius of the RESONANT orbit of an electron in a hydrogen atom. But the obtained value of the radius of the orbit is approximately 4 times larger than the radius of the hydrogen atom obtained from the experiment.

If we assume that the frequency of a light quantum is equal to HALF the frequency of rotation of an electron in orbit, then instead of equations (9-10) we get

$$\frac{\hbar\omega_1}{2} = m \frac{\omega_1^2 r_1^2}{2} \Rightarrow \omega_1 = \frac{\hbar}{mr_1^2} \quad (13)$$

And instead of equation (11) we have

$$\frac{\hbar}{mr_1^2} = \frac{e}{2\sqrt{\pi\epsilon_0 m}} \frac{1}{r_1^{3/2}} \quad (14)$$

And, then, we obtain the expression of the radius of the orbit determined by the World Constants, which gives a value that coincides with the experimental value of the hydrogen radius given under Figure 1 with good accuracy.

$$r_1 = \frac{1}{4} \frac{16\hbar^2\pi\epsilon_0}{e^2 m} = 53.03\text{pm} \quad (15)$$

At the same time, taking into account that the particle is an electron, it is a packet of waves, it is natural to assume that the resonant orbit corresponds to a resonant wave. Plasma, resonant, i.e. standing fluctuations in the charge density lead to the formation of charged layers at the boundaries of the resonator. But there are no boundaries in a circular orbit and, consequently, there are no areas of charge accumulation - there is no dipole. Therefore, again, it is natural that this resonance wave does not give a charge to the radiation of an electromagnetic wave. And the dipole

arises only between the entirely displaced orbit and the nucleus. So the excitation of the entire electron orbit is parametric, as it should be at half the frequency of its intra-orbital resonance!

So, in fact, the rejection of the Bohr orbital model of the atom was incorrectly "justified" by the instability of the (energy) of this electron orbit due to electromagnetic radiation. But, taking into account the corpuscular-wave dualism, the de Broglie wave packet of an electron in a resonant orbit also enters into resonance - in a self-coherent state, i.e. the electron becomes a resonant undammed wave, which, as noted above, does not have a dipole and does not radiate at the resonant frequency.

And taking into account the obtained parametric relationship between the frequencies of rotation of an electron in a resonant orbit and the quantum of light exciting it from the orbit, one can also imagine the parametric relationship of the de Broglie wave with an electromagnetic wave. In this case, the de Broglie wavelength of an electron in a resonant orbit is determined by the circumference of this orbit

$$\lambda_{dB} = 2\pi r_1 = \frac{8\hbar^2 \pi^2 \epsilon_0}{e^2 m} \quad (16)$$

So the ideal (undamped) resonance of the de Broglie wave DOES NOT radiate energy, if the Planck quantum for electromagnetic waves is extended not only to acoustic (as Einstein did), but also to the de Broglie waves of the electron.

$$v_2^2 = \frac{e^2}{2\pi\epsilon_0 r_2 m} \Rightarrow v_2 = \pm \frac{e}{\sqrt{2\pi\epsilon_0 m r_2}} = \pm 3978566.4 \text{ m/s} = \pm 0.0133 \cdot c \quad (19)$$

Wherein again, the classical consideration gives the first functional relationship between the speed of a pair of electrons in an orbit and its radius and the total potential energy of this pair of electrons.

$$U_{22} = \frac{e^2}{\pi\epsilon_0 r_2} = 179.548356 \text{ eV} \quad (20)$$

Where one electron accounts for half of the potential energy of the pair:

$$U_{21} = \frac{U_{22}}{2} = \frac{e^2}{2\pi\epsilon_0 r_2} = 89,7742 \text{ eV} \quad (21)$$

At a higher speed in the orbit of a pair of electrons from formula (19), the kinetic energy of a pair of electrons in this orbit naturally also increases, but again it is exactly half of the potential energy of a pair of electrons:

$$E_c = 2m \frac{v_2^2}{2} = 2m \frac{2\pi\epsilon_0 r_2 m}{2} = \frac{e^2}{2\pi r_2 \epsilon_0} = U_{21} = I_{22} \quad (22)$$

So the energy diagram for a pair of electrons, and for each electron of this pair, is completely similar to the diagram

### Principles of formation of multi-electron orbitals

The representation of an electron orbital as a packet of de Broglie waves of an electron transformed into a resonant wave gives a simple explanation for the Pauli principle. Given that orthogonally polarized (charge) waves do not repel each other, it can be assumed that two electron waves can simultaneously be in the same orbit as one particle with doubled mass and doubled charge. This corresponds to the Pauli principle for electrons as particles. Therefore, for this pair of electrons in orbit, the centrifugal force, in contrast to formula (1), is equal to

$$F_C = 2m\omega^2 r = 2m \frac{v^2}{r} \quad (17)$$

And the force of the Coulomb attraction of a pair of electrons in this orbit to a nucleus with a double charge (helium atom) is equal to

$$F_{22} = \frac{e^2}{\pi\epsilon_0 r^2} \quad (18)$$

From the equality of forces in the orbit of this pair of electrons, similarly to formula (3), but now using the radius of the helium atom  $32 \cdot 10^{-12} \text{ m}$ , we obtain a large, but also quite reasonable speed of movement of this pair of electrons along the orbit of the order of a percent from the speed of light.

Shown in Fig 2. And to pull out a pair of electrons from this level, you need eV, which is close to the Total Ionization Energy attributed to helium  $-79.005151042(40) \text{ eV}$  and in order to pull out one electron from this orbit, half of this Ionization Energy of the pair is required -  $44.8871 \text{ eV}$ . But this energy value is almost twice the tabular value of the First Ionization Potential of helium  $24.47 \text{ eV}$ .

It also follows from the above reasoning that the level of the remaining one electron of the helium atom decreases after primary ionization, and during ionization it needs to additionally overcome the Coulomb force of the nucleus charged by the first ionization. But for the time being, we will not be distracted by these arguments, which may and will help explain the discrepancies in the energy levels of the helium atom, which is more complex than hydrogen. And immediately we will use the quantum addition to the classical calculations of the orbital of a pair of electrons on the basis of its energy diagram obtained.

To do this, first again, from the equality of forces acting on a pair of electrons in orbit, in the classical consideration, we can obtain, similarly to formula (8), in addition to (19), the dependence of the frequency of rotation of this pair around the nucleus on the radius of the pair orbit

$$2m\omega^2 r = \frac{e^2}{\pi\epsilon_0 r^2} \Rightarrow \omega_2^2 = \frac{e^2}{2\pi\epsilon_0 r_2^3 m} \Rightarrow \omega_2 = \frac{e}{\sqrt{2\pi\epsilon_0 m}} \frac{1}{r_2^{3/2}} \quad (23)$$

We use the Planck equation and the Planck constant to relate to the frequency of light, the quantum of which is equal to the energy of the Ionization Potential and which, according to formula (22), is equal to the kinetic energy of a pair of electrons in orbit

$$I_{22} = \hbar \omega_{ph2} = E_{C2} = 2m \frac{v_2^2}{2} = m \omega_2^2 r_2^2 \quad (24)$$

This equation complements the classical ones and, thus, gives an additional connection between the rotation frequency and the radius of the orbit. And, thus, if we ASSUME that the frequency of a light quantum correlates with the frequency of light, it makes it possible to obtain a single value of both the frequency and the radius of the classical orbit.

In particular, if these frequencies are equal, then we obtain, in addition to equation 8, another equation relating the frequency to the radius.

$$\omega_2 = \frac{\hbar}{mr_2^2} \quad (25)$$

And equating the frequencies, we again get one equation with one unknown radius

$$\frac{\hbar}{mr_2^2} = \frac{e}{\sqrt{2\pi\epsilon_0 m} r_2^{3/2}} \quad (26)$$

The solution of the obtained equation gives us the only value of the radius of the RESONANT orbit of a pair of electrons in a helium atom, which is 8 times less than the radius of an electron orbit in hydrogen.

$$\frac{\hbar}{e} \sqrt{\frac{2\pi\epsilon_0}{m}} = r_2^{1/2} \Rightarrow r_2 = \frac{2\pi\epsilon_0 \hbar^2}{me^2} = \frac{1}{8} r_1 = 26.52 \text{ pm} \quad (27)$$

This one, obtained from the World Constants, with equal frequencies of light and rotation of a pair of electrons per orbit,  $e$ , the radius of the circular orbit of rotation of a pair of electrons 26.52 pm is close to the radius of helium 32 pm. Consequently, the parametric coupling is weaker, and, as already noted, the dipole of electron waves does not appear on the orbit. Therefore, the excitation of the entire pair of electrons is unlikely. But we will not be special consideration of the Second Ionization Potentials in this work. We simply note that the use of the quasi-nuclear model for a completely filled shell of a pair of electrons makes it possible, even with an increase in the atomic number, to estimate from the first principles both the radius of the additional electron shell and the energy level corresponding to it, as approximately equal to their values in the hydrogen atom. This numerically differs from the experimental values by tens of percent (Fig. 4, blue dotted

lines), which is natural in the first approximation by the quasi-nuclear model, which does not take into account the repulsion of the upper electron by the filled inner shell. But even these discrepancies, which can be seen how to eliminate, are not catastrophic, by orders of magnitude, which gives the Schrödinger equation [4, 12].

The quasi-nuclear model with the calculation algorithm described above gives a significant jump in the size of the atom and its First Ionization Potential even when one more electron is added to the completely filled named c-shell. And here there is a relatively small change in the atomic number and radius of the orbit of the added outer electron, and its First Ionization Potential (Fig. 4, red dotted lines). But for a more rigorous calculation than the quasi-nuclear model, according to the described algorithm, of course, it is necessary to take into account the symmetry of the tetrahedron of electron shells when the c-shell is completely filled for the second and third periods, and their even more complex symmetry for the following periods, which manifests itself during the transition from - shells to the so-called p-shell (green dotted curves in Fig. 4), where the equivalent orbitals form, as previously shown, a tetrahedron.


And so, as it should be, correctly chosen Basic Models give/describe the correct order of observed phenomena, in contrast to erroneous, but Canonized Models, which give catastrophic discrepancies with experiment and are suitable only for decoration and deception of the layman.

And the algorithm described in this work allows us to obtain from the First Principles the "Second" Principle - the Principle of the formation of multi-electron orbitals and the corresponding levels of allowed (Resonance) energies in an atom with any atomic number.

The deepest orbital and its energy level, the deepest, are formed in the Coulomb field of the nucleus, a field from the maximum number of equivalent orbitals satisfying point spatial symmetry (this number of electrons of the inner shell is less than or equal to the total number of electrons of the atom).

The remainder of the total number of electrons, in turn, is divided by the maximum number of electrons for reduced symmetry, which form the next higher allowed energy level.

Thus, in a recurrent way, using the quasi-nuclear model, it is possible to calculate from the First Principles (from the World Constants) - from bottom to top, all the electronic levels of different atoms, from an atom with a completely filled electron shell to atoms filling with electrons a new shell with a quasi-nucleus from the nucleus and the last completely filled shells. This approach gives the correct order of allowed electronic levels, corresponding to the Ionization Potentials. Only for a quantitatively rigorous calculation it is necessary to take into account the degree of shielding of the nucleus by deeper filled shells, which, naturally, is higher for the lowest shell with the maximum number of symmetry degeneracy, but is enhanced by subsequent-overlying ones and is minimal for the upper shell of one orbit with two electrons.



**Fig 4:** Experimental dependences of the radius of the outer electron shell and its First Ionization Potential on the atomic number (point) and their assessment based on the Fundamental Constants according to the Principles described in the work (dashed lines). Zero denotes the “zero” element used in the construction of the periodic table by Mendeleev, and minus one is an additional conditional element with minus one electron.

## References

- Born M. Atomic physics, 2nd ed. - Moscow: Publishing House World; c1967. p. 493.
- Qian Xue-Sen. Physical Mechanics, Moscow: Publishing House World; c1965. p. 427.
- Ordin SV. « CHAOS – Imaginary Ostensibility – Orthogonality », Global Journal of Science Frontier Research - Physics & Space Science, GJSFR-A. Version 1.0. 19(3):49-58, [https://globaljournals.org/GJSFR\\_Volume19/3-Chaos-Imaginary-Ostensibility.pdf](https://globaljournals.org/GJSFR_Volume19/3-Chaos-Imaginary-Ostensibility.pdf)
- Ian M, Tomislav C, Klays H, Nikola K, Kozo K, Quantities, Units and Symbols in Physical Chemistry, Second Edition, Blackwell Science, IUPAC Gold Book. Retrieved 2012 April 3. Archived from the original on 2012 March 2.
- Ordin SV. Frontier Chemistry Aspects, Global Journal of Science Frontier Research: B-Chemistry (GJSFR-B). Version 1.0. 2020;20(2):1-11. ISSN Online: 2249-4626, ISSN Print: 0975-5896, DOI: 10.17406/GJSFR, [https://globaljournals.org/GJSFR\\_Volume20/1-Frontier-Chemistry-Aspects.pdf](https://globaljournals.org/GJSFR_Volume20/1-Frontier-Chemistry-Aspects.pdf)
- Marat VG. Wave Function is not amplitude of Probability Density. American Journal of Modern Physics. 2017;6(4):49-50. DOI: 10.11648/j.ajmp.20170604.11
- Ordin SV. « Newton's Coulomb Laws », Global Journal of Science Frontier Research- Physics & Space Science (GJSFR-A). Version 1.0, 2019;19(1):145-155. [https://globaljournals.org/GJSFR\\_Volume19/E-Journal\\_GJSFR\\_\(A\)\\_Vol\\_19\\_Issue\\_1.pdf](https://globaljournals.org/GJSFR_Volume19/E-Journal_GJSFR_(A)_Vol_19_Issue_1.pdf)
- Stanislav O. Review Article, Non-Elementary Elementary Harmonic Oscillator, American Journal of Materials & Applied Science (AJMAS). Volume 3 Issue 1, Published Date: 03/08/2021, Pages: 003-008, <https://www.scireslit.com/MaterialScience/AJMAS-ID14.pdf> <https://www.scireslit.com/MaterialScience/articles.php?volume=3&issue=1>



9. Stanislav VO. Book: Foundations Of Planck-Einstein Quantization (Thematic Collection of Recent Studies Reviewed in Scientific Journals), LAP Lambert Academic Publishing, ISBN 978-620-4-21066-7; c2021. p. 110.
10. Ordin SV. « Parametrically excited Anharmonic Oscillator », Global Journal of Science Frontier Research - Physics & Space Science, GJSFR-A. Volume 19 Issue 3. Version 1.0 p. 133-144.  
[https://globaljournals.org/GJSFR\\_Volume19/7-Parametrically-Excited.pdf](https://globaljournals.org/GJSFR_Volume19/7-Parametrically-Excited.pdf)
11. Stanislav O. Book: Refinement of basic physical models, Lambert, Project No 163273, ISBN: 978-3-659-86149-9; c2017. p. 82.
12. Stanislav O. Letter to Editor, Evidence-Based Physics, American Journal of Materials & Applied Science. 3(1):001-002. Published Date: 02/22/2021.  
<https://www.scireslit.com/MaterialScience/AJMAS-ID13.pdf>
13. Stanislav O. Gaps and Errors of the Schrödinger Equation, Global Journal of Science Frontier Research, GJSFR. Version 1.0, 22(3):1-5.  
[https://globaljournals.org/GJSFR\\_Volume22/3-Gaps-and-Errors.pdf](https://globaljournals.org/GJSFR_Volume22/3-Gaps-and-Errors.pdf)
14. Stanislav O. Analysis of Newton's Elementary Particle, International Journal of Recent Scientific Research, IJRSR –18424/2021
15. Stanislav O. « Dualism of newton's elementary particle », International Journal of Physics and Applications (IJOS), E-ISSN: 2664-7583, P-ISSN: 2664-7575, IJOS. 2022;4(1):07-16.
16. Stanislav O. « Imagination and real quantization », International Journal of Physics and Applications, E-ISSN: 2664-7583, P-ISSN: 2664-7575, IJOS. 2022;4(1):35-38.
17. Ordin SV. Quasinuclear foundation for the expansion of quantum mechanics, International Journal of Advanced Research in Physical Science (IJARPS). 2018;5(6):35-45, Paper ID # 3115.  
<https://www.arcjournals.org/international-journal-of-advanced-research-in-physical-science/volume-5-issue-6>